

12.2 Linear momentum, angular momentum, work and energy

If you know a particle's starting position and velocity, and you know the force on it as it moves, then you can use $\vec{F} = m\vec{a}$ to predict its path. That is the central idea of the previous section. We had no need for ideas related to momentum, angular momentum, work and energy. For one particle the one equation $\vec{F} = m\vec{a}$ tells the whole story. So, before we go on to discuss them further, let us be clear:

The concepts of linear and angular momentum, work and energy are not needed to study particle mechanics. $\vec{F} = m\vec{a}$ is enough.

So, why do we bother to devote a section to these topics? Because

- These concepts will sometimes be needed when we discuss more complex systems;
- These concepts sometimes provide a shorter route for answering some dynamics questions;
- The simplest place to introduce the concepts is in the context of one particle;
- The concepts give a way to check the consistency of solutions of $\vec{F} = m\vec{a}$; and
- The concepts can be an aid to physical intuition.

For more complex systems, principles of momentum and energy transcend $\vec{F} = m\vec{a}$ and can generally not be derived from $\vec{F} = m\vec{a}$. But for a single particle, all of these are derived concepts, as worked out in box 12.4 on page 9. Note that

All of the facts and theorems below apply to any motion of a particle that is consistent with $\vec{F} = m\vec{a}$.

Example: Simple ballistics solution.

Consider a ball thrown up at 45°: $\vec{F} = -mg\hat{j}$, $\vec{r}(0) = \vec{0}$ and $\vec{v}(0) = v_0\hat{i} + v_0\hat{j}$. We claimed (page 616) that a solution is

$$\vec{r} = v_0 t \hat{i} + (v_0 t - gt^2/2) \hat{j} \quad \text{and} \quad \vec{v} = v_0 \hat{i} + (v_0 - gt) \hat{j}.$$

This solution is plotted various ways in fig. 12.22. These functions of time are consistent with the initial conditions. Further they are consistent with the governing equations, the so called 'equations of motion', $\vec{v} = \vec{F}/m$ and $\vec{r} = \vec{v}$. All of the momentum and energy principles below must therefore apply.

Some of the ideas apply even if $\vec{F} \neq m\vec{a}$. For example, the work of a force is defined for imagined motions that might never occur.




Figure 12.22: (a) A simple ballistics problem. (b) The well known parabolic trajectory. (c) The velocity trajectory, called a *hodograph*, shows the sequence of positions of the tip of the velocity vector. The velocity starts with $v_x = v_y$ and then v_y decreases. (d) Plot of x and y components vs time. The y component has a parabolic shape similar to the particle trajectory because x is proportional to t in this problem. (e) The velocity components versus time. Note, (b) is a 'cross plot' of the two plots in (d), and (c) is a cross plot of (e).

Linear momentum

Linear momentum for a particle is defined as $\vec{L} = m\vec{v}$. The particle momentum-balance theorems (facts) are

$$\vec{F} = \frac{d}{dt}\vec{L} \quad \text{and} \quad \underbrace{\int_{t_1}^{t_2} \vec{F} dt}_{\text{Linear impulse}} = \vec{L}_2 - \vec{L}_1$$

These are so trivially related to $\vec{F} = m\vec{a}$ that it is hard to see any content in them. And, indeed, if we were only studying the mechanics of single particles we probably would not have introduced the concept of linear momentum. Nonetheless, the general result does apply:

The net force $\vec{F}$ on a particle is the rate of change of its linear momentum, $\dot{\vec{L}}$.

A special important case is when there is no force and linear momentum is conserved (doesn't change). For a single particle momentum conservation means constant velocity motion.

Example: Linear momentum check

For the simple ballistics solution above we evaluate the left side of the momentum balance equation

$$\int_0^t \vec{F} dt' = -mgt\mathbf{j}.$$

Then evaluate the right side:

$$\Delta\vec{L} = \vec{L}_2 - \vec{L}_1 = [m(v_0\mathbf{i} + (v_0 - gt)\mathbf{j})] - [m(v_0\mathbf{i} + v_0\mathbf{j})] = -mgt\mathbf{j}$$

and check for equality: $-mgt\mathbf{j} = -mgt\mathbf{j}$. This force and momentum is plotted in fig. 12.23. The solution is consistent with linear momentum balance. Note that in this example there is no change in the component of linear momentum in the t direction; there is no force in the x direction so L_x is conserved.




Figure 12.23: The force and linear momentum from the problem in fig. 12.22. Note that the momentum curves are the integrals of the corresponding force curves.

Angular momentum

In dynamics angular momentum is one of the most important ideas, important theoretically and for solving problems. For one particle

Angular momentum relative to point C is $\vec{H}_C = \vec{r}_C \times (m\vec{v})$,

where $\vec{r}_C$ is the position of the particle relative to fixed point C and $\vec{v}$ is the velocity of the particle. Angular momentum can be calculated relative to any point C. Which point you pick affects the value of the angular momentum. Sometimes $\vec{H}_C$ is written without the “/” as $\vec{H}_C$. The key angular momentum theorems (facts) are:

$$\underbrace{\vec{r}_{/C} \times \vec{F}}_{\vec{M}_C} = \dot{\vec{H}}_{/C} \quad \text{and} \quad \underbrace{\int_{t_1}^{t_2} \vec{r}_{/C} \times \vec{F} dt}_{\text{Angular impulse}} = (\vec{H}_{/C})_2 - (\vec{H}_{/C})_1.$$

The torque $\vec{M}_C$ of all the external forces acting on a particle about point C is the rate of change of its angular momentum $\dot{\vec{H}}_{/C}$ about point C.

The intuitive notion is that angular momentum represents how much a particle is ‘going around’ point C. A particle gets more credit for going faster, for being more massive, and for being farther away^①.

If the force on the particle is zero or passes through the point C, the torque (moment) of the force is zero and its angular momentum is conserved.

Example: Angular momentum check.

Using the same ballistics example we check the solution for consistency with angular momentum balance. For no good reason let’s use the origin for the angular momentum reference point. We could use any point. Again we compare the left and right sides and check for equality.

$$\begin{aligned} \underbrace{\int_0^t \vec{r}_{/0} \times \vec{F} dt}_{\text{Left side}} &= \int_0^t (v_0 t' \hat{i} + (v_0 - gt'^2/2) \hat{j}) \times (-mg \hat{j}) dt' \\ &= \int_0^t -v_0 m g t' \hat{k} dt' = -v_0 m g \hat{k} t^2/2 \\ \underbrace{\vec{H}_{/02} - \vec{H}_{/01}}_{\text{Right side}} &= \vec{r}(t) \times (m \vec{v}(t)) - \vec{r}(0) \times (m \vec{v}(0)) \\ &= m (v_0 t \hat{i} + (v_0 t - gt^2/2) \hat{j}) \times (v_0 \hat{i} + (v_0 - gt) \hat{j}) - m (\vec{0} \times \vec{v}(0)) \\ &= (v_0 t - gt^2 - v_0 t + gt^2/2) m v_0 \hat{k} = -v_0 m g t^2 \hat{k} / 2. \end{aligned}$$

This torque and angular momentum are plotted in fig. 12.24. The ‘left side’ agrees with the ‘right side’ and angular momentum balance is satisfied, as it must be. In 2D problems like this there is only one non-trivial component to angular momentum, that in the out-of-plane (z) direction. In this case the angular momentum H_z is not conserved because the gravity force does have a torque about the z axis.

Power balance

The power P of a force $\vec{F}$ on a particle with velocity $\vec{v}$ is $P = \vec{F} \cdot \vec{v}$. The kinetic energy of a particle is $E_K = mv^2/2$. The power balance equations are

$$P = \dot{E}_K \quad \text{and} \quad \int_{t_1}^{t_2} P dt = E_{K2} - E_{K1}.$$

① **Angular momentum contradicts common intuition.** The notion that angular momentum is bigger if a given mass is further away is counterintuitive to most of us. Imagine a one kilogram mass is going around your head once per second at a distance of a meter. Now imagine a second equal mass going around at a radius of 10 meters and only going around once every 100 seconds. The first mass whirls around your head 100 times for each slow orbit of the second mass. Even though the second mass certainly doesn’t make its rotation feel so present its angular momentum is as big. Intuitive or not, this is how angular momentum is defined. It’s a useful concept so it’s worth adjusting your intuition to match it.




Figure 12.24: The torque and angular momentum about the origin for the problem in fig. 12.22. For this 2D problem only the z component is non-trivial. Note that the angular momentum is the integral of the torque.

The power P of *all* the external forces acting on a particle is the rate of change of its kinetic energy $\dot{E}_K$. Or, integrating in time, the power added up over time is the net change in kinetic energy.

The situation is similar to that for 1D motion (section 10.2).

Power and work

The integral of power with respect to time can be replaced with a path integral for the work of a force. The key idea is in the differential expressions for an increment of work:

$$dW = P dt = \vec{F} \cdot \vec{v} dt = \vec{F} \cdot \frac{d\vec{r}}{dt} dt = \vec{F} \cdot d\vec{r}.$$

So

$$\begin{aligned} W_{12} &= W_{12} \\ \text{Power integrated in time} &= \text{Sum of work increments} \\ \int_{t_1}^{t_2} P dt &= \int_1^2 dW \\ \int_{t_1}^{t_2} \vec{F} \cdot \vec{v} dt &= \int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot d\vec{r} \end{aligned} \quad (12.8)$$

Thus the power balance equation, integrated in time is equivalent to the “work energy” equation:

$$\begin{aligned} \text{Work on particle} &= \text{Change in its kinetic energy} \\ W_{12} &= \Delta E_K \\ &= E_{K2} - E_{K1} \end{aligned} \quad (12.9)$$

Example: Energy check.

We can check the simple ballistics solution for consistency with energy balance.

First let's compare the work to the change of kinetic energy.

$$\begin{aligned}
 \underbrace{\int_0^t \vec{F} \cdot \vec{v} dt}_{\text{Work}} &= \int_0^t (-mg\hat{j}) \cdot (v_0\hat{i} + (v_0 - gt')\hat{j}) dt' \\
 &= \int_0^t -mg(v_0 - gt') dt' = -mgv_0t + mg^2t^2/2 \\
 \underbrace{E_{K2} - E_{K1}}_{\text{Change in kinetic energy}} &= m(|\vec{v}|^2 - |\vec{v}_0|^2)/2 \\
 &= m((v_0^2 + (v_0 - gt)^2 - 2v_0^2)/2) = \underbrace{-mgv_0t + mg^2t^2/2}_{\text{agrees}}
 \end{aligned}$$

This power and kinetic energy are plotted in fig. 12.25. In this check we have not taken advantage of the fact that this particular force is conservative.

The work of a force $\vec{F}$: W_{12}

Previously in Physics, and more recently in one dimensional dynamics here, you learned that

Work is force times distance.

This is actually a special case of the formula

$$P = \vec{F} \cdot \vec{v}.$$

How is that? If $\vec{F}$ is constant and parallel to the displacement $\Delta\vec{x}$, then

$$\begin{aligned}
 W_{12} &= \int \dot{W} dt = \int P dt = \int \vec{F} \cdot \underbrace{\vec{v} dt}_{d\vec{x}} = \int \vec{F} \cdot d\vec{x} = \vec{F} \cdot \int d\vec{x} \\
 &= \vec{F} \cdot \Delta\vec{x} = F\Delta x = \text{Force} \cdot \text{distance}.
 \end{aligned}$$

In 2 and 3 dimensions there are subtleties involved with the concept of work because of its dependence on which path in space the force works on. These 'path dependence' subtleties are often covered in some detail in calculus courses in the sections on vector calculus, path integrals, gradient and curl. We discuss the relevant highlights below.

Potential energy of a force

Some forces (read force fields $\vec{F} = \vec{F}(\vec{r})$) have the property that the work they do is independent of the path followed by the material point as the force acts. If the work of a force is path independent in this way (see box 12.3 on page 7), then a potential energy can be defined so that the work done by the force is the decrease in the Potential Energy




Figure 12.25: The power of the gravitational force and the total kinetic energy are plotted vs time for the ballistics problem of fig. 12.22. Note that the kinetic energy is the integral of the power. The gravitational power starts out negative, goes to zero when the particle reaches the apex of its trajectory and then becomes positive evermore as the downwards gravitational force acts on a downwards moving particle. The kinetic energy starts at mv_0^2 and drops to $mv_0^2/2$ at the particle apex when v_y goes to zero. Then the kinetic energy increases forever more as the gravitational force does more and more work.

$$-\Delta E_P = W_{12} = E_{P1} - E_{P2}$$

The common examples are listed below:

- **linear spring:** $E_P = (1/2)k(\text{stretch})^2$.
- **gravity near earth's surface:** $E_P = mgh$
- **gravity between spheres or points:** $E_P = -MmG/r$
- **constant force $\vec{F}$ acting on a point:** $E_P = -\vec{F} \cdot \vec{r}$

In all cases a constant could be added to the potential energy and it would still be a legitimate potential energy for the force.

In the cases of the spring and gravity between spheres, the change in potential energy is the net work done by the spring or gravity on the pair of objects between which the force acts. If both ends of a spring are moving, the net work of the spring on the two objects to which it is connected is the decrease in potential energy of the spring.

There is a possible source of confusion in our using the same symbol E_P to represent the potential work of an external force and for internal potential energy. In practice, however, they are used identically, so we use the same symbol for both. The potential energy in a stretched spring is the same whether it is the cause of force on a system or it is internal to the system.

Example: Checking conservation of energy

Because the gravity force is conservative we can also check our simple ballistics solution for consistency with conservation of energy. Taking the potential energy as $E_P = mgh = mgy$ we find, as expected, that the solution does have the property that

$$\begin{aligned} E_{tot1} &\stackrel{?}{=} E_{tot2} \\ E_{K1} + E_{P1} &\stackrel{?}{=} E_{K2} + E_{P2} \\ mv_0^2 + 0 &\stackrel{?}{=} (v_0^2 + (v_0 - gt)^2)m/2 + mg \overbrace{(v_0 t - gt^2/2)}^y \\ mv_0^2 &\stackrel{\checkmark}{=} mv_0^2 \quad (\text{Checks}) \end{aligned}$$

Using momentum and energy as a check of a numerical solution

You obtain a numerical solution to $\vec{F} = m\vec{a}$ by setting up the set of first order differential equations 12.2 on page 615. In turn, these can be written in explicit scalar form as eqn. (12.3).

While you solve these equations you can add further first order equations that you can use in your energy and momentum checks. These eval-

uate the integrals for linear impulse, angular impulse and work.

$$\begin{aligned}\frac{d}{dt}(\text{linear impulse}) &= \vec{F}, \\ \frac{d}{dt}(\text{angular impulse}) &= \vec{r}_{/C} \times \vec{F}, \text{ and} \\ \dot{W} &= \vec{F} \cdot \vec{v}.\end{aligned}$$

The first two equations are short hand for 2 (or 3) first order scalar equations for motion in 2 (or 3) spatial dimensions. If these are added to the system of ordinary differential equations that you solve, they can be used to check the solution.

Summary on using energy and momentum to check a solution

Because the momentum and energy facts and theorems apply to any motion consistent with $\vec{F} = m\vec{a}$ they can be used as a consistency check on any solutions you find to the differential equations of motion ($\vec{F} = m\vec{a}$).

Here is the general situation. You are given $\vec{F}(t, \vec{r}, \vec{v})$. You are given initial conditions $\vec{r}_0$ and $\vec{v}_0$ at, say, $t = 0$. Using computer integration or pencil and paper methods, you solve the differential equation $\vec{a} = \vec{F}/m$ to get $\vec{r}(t)$ and $\vec{v}(t)$. Now your solution can be checked for consistency with the energy and momentum theorems. In particular, your solution, if it is correct, must satisfy

- Linear momentum balance: $\int_0^t \vec{F} dt = \vec{L}_2 - \vec{L}_1$;
- Angular momentum balance: $\int_0^t \vec{r} \times \vec{F} dt = \vec{H}_2 - \vec{H}_1$;
- Work-energy: $\int_0^t \vec{F} \cdot \vec{v} dt = E_{K2} - E_{K1}$.

12.3 Conservative forces and non-conservative forces

Imagine that the force $\vec{F}$ on a particle is known to depend on the position $\vec{r}$ of a particle as it moves. This dependence of $\vec{F}$ on $\vec{r}$ is called a *force field*:

$$\vec{F} = \vec{F}(\vec{r}).$$

As the particle moves from one point $\vec{r}_1$ to another $\vec{r}_2$ we can evaluate the work of this force field as

$$W_{12} = \int_{t_1}^{t_2} P dt = \int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot \vec{v} dt = \int_{\vec{r}_1}^{\vec{r}_2} \vec{F}(\vec{r}) \cdot d\vec{r}.$$

But what if the particle moves between the same two points but along a different path, is the work W_{12} the same? If that is true then the work going from $\vec{r}_1$ to $\vec{r}_2$ and back would be zero. Which means the work of the force when the particle moves on any closed path would be zero. Here is an example of a force field $\vec{F}(\vec{r})$ in the xy plane where the work in going on a closed path, from $\vec{r}_1$ to $\vec{r}_1$, from home to home, is not zero:

$$\vec{F} = C[\hat{k} \times \vec{r}] = C[-y\hat{i} + x\hat{j}]$$

where C is a constant. This force pushes the particle around in circles. So, if the particle moves on the circular path

$$\vec{r} = r_0[\cos\theta\hat{i} + \sin\theta\hat{j}] \quad (0 \leq \theta \leq 2\pi)$$

then the work is the force magnitude times the arc-length (the force is parallel to the velocity for this path) and so,

$$\int_{\theta_1}^{\theta_2} \vec{F} \cdot d\vec{r} = 2\pi C r_0^2 \neq 0.$$

This force field gives non-zero work for some closed paths, thus is path dependent for open paths and therefore is *non-conservative*. How can you tell if a force field is conservative or not. This, you learn in vector calculus, holds if the curl of $\vec{F}$ is zero, $\vec{\nabla} \times \vec{F} = \vec{0}$, everywhere.

Forces from any combination of springs and gravity are always conservative.

These have been used in the simple ballistics example above. That linear momentum balance, angular momentum balance and energy balance, all are consistent to an assumed solution lends credence to its correctness. For simple problems with such simple analytical solutions, using this consistency is not the most efficient way of checking a candidate solution's veracity. We would be better off just plugging the proposed solution back into the differential equation to see if it was satisfied. But in more complex problems and in numerical solutions, checks like those here are sometimes simpler to make. Some more comments about these checks:

- The angular momentum check can be used relative to any fixed point you choose. If you can find a point where, say, the applied force has no moment, then the change of angular momentum should be zero about that point.
- If the applied force is conservative, the work integral can be replaced by the change in potential energy and the work-energy check is a check of the conservation of energy.
- If you try to make the checks with pencil and paper the checks can sometimes be harder to implement than it was to find the original solution.
- These checks are often *very useful*, and this is perhaps an understatement, for checking the validity of numerical solutions of dynamics equations. Basically you shouldn't trust yours or any body else's code unless such checks have been made. It is hard to write correct code without making such checks. And such checks are a strong sign of code reliability because an error in computer code will usually lead to an error in momentum balance, angular momentum balance or energy balance.

12.4 Derivation of momentum, angular momentum and energy theorems for a point mass

For a point-mass particle the principles of linear momentum, angular momentum and energy are theorems that can be derived simply from

$$\vec{F} = m\vec{a}$$

as follows.

Linear momentum

Define linear momentum as $\vec{L} = m\vec{v}$ then differentiating we have the equation $\vec{F} = m\vec{a}$. It is not so much a derivation but a restatement to write:

$$\vec{F} = \dot{\vec{L}}$$

Integrating both sides in time we get

$$\underbrace{\int_{t_1}^{t_2} \vec{F} dt}_{\text{impulse}} = \underbrace{\vec{L}_2 - \vec{L}_1}_{\text{change of momentum}}.$$

This is the principle of impulse and momentum.

Angular momentum

Start with $\vec{F} = m\vec{a}$ and take the cross product of both sides with the position relative to a fixed point C and you get

$$\vec{r}_{/C} \times \vec{F} = \vec{r}_{/C} \times (m\vec{a}).$$

Now if we define $\vec{H}_{/C} = \vec{r}_{/C} \times (m\vec{v})$ we can differentiate to find that, writing out all details,

$$\begin{aligned} \dot{\vec{H}}_{/C} &= \frac{d}{dt} (\vec{r}_{/C} \times (m\vec{v})) \\ &= \frac{d}{dt} ((\vec{r} - \vec{r}_C) \times m\vec{v}) \\ &= m(\dot{\vec{r}} - \dot{\vec{r}}_C) \times \vec{v} + m(\vec{r} - \vec{r}_C) \times \dot{\vec{v}} \\ &= m(\vec{v} - \vec{0}) \times \vec{v} + m(\vec{r} - \vec{r}_C) \times \dot{\vec{v}} \\ &= \underbrace{m\vec{v} \times \vec{v}}_{\vec{0}} + m(\vec{r} - \vec{r}_C) \times \dot{\vec{v}} \\ &= \vec{r}_{/C} \times (m\dot{\vec{v}}) \\ &= \vec{r}_{/C} \times (m\vec{a}) \end{aligned}$$

Putting these together we have

$$\vec{r}_{/C} \times \vec{F} = \dot{\vec{H}}_{/C}.$$

Integrating both sides with respect to time we get that the net angular impulse is the change in angular momentum.

$$\int_{t_1}^{t_2} \vec{r}_{/C} \times \vec{F} dt = (\vec{H}_{/C})_2 - (\vec{H}_{/C})_1.$$

Power and kinetic energy

The power equation is found with a shade more difficulty. We take the equation $\vec{F} = m\vec{a}$ and dot both sides with the velocity $\vec{v}$ of the particle:

$$\vec{F} \cdot \vec{v} = m\vec{a} \cdot \vec{v}. \quad (12.10)$$

Evaluating $\vec{v} \cdot \vec{a}$ is most easily done with the benefit of hindsight. So we cheat and look at the time derivative of the speed squared:

$$\begin{aligned} \frac{d}{dt} \left(\frac{1}{2} v^2 \right) &= \frac{1}{2} \frac{d}{dt} (\vec{v} \cdot \vec{v}) \\ &= \frac{1}{2} (\dot{\vec{v}} \cdot \vec{v} + \vec{v} \cdot \dot{\vec{v}}) \\ &= \vec{v} \cdot \dot{\vec{v}} \\ &= \vec{v} \cdot \vec{a} \end{aligned}$$

Applying this result to eqn. (12.10) we get

$$\underbrace{\vec{F} \cdot \vec{v}}_P = \frac{d}{dt} \left(\underbrace{\frac{1}{2} m v^2}_{E_K} \right),$$

the energy (or power balance) equation for a particle.

Integrating in time we get

$$\underbrace{\int_{t_1}^{t_2} \vec{F} \cdot \vec{v} dt}_{\text{Work}} = \underbrace{E_{K2} - E_{K1}}_{\text{Change in kinetic energy}},$$

Power and work and energy

Because $\vec{F} \cdot \vec{v} dt = \vec{F} \cdot d\vec{r}$ the time integral of power can be replaced with a path integral, the standard work integral:

$$\int_{t_1}^{t_2} \vec{F} \cdot \vec{v} dt = \int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot d\vec{r}.$$

If $\vec{F}$ is a conservative force field, meaning a function of position, then $E_P(\vec{r})$ exists, so that

$$-\vec{\nabla} E_P = \vec{F} \quad \text{then} \quad \int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot d\vec{r} = E_P(\vec{r}_1) - E_P(\vec{r}_2)$$

and the work-energy equation becomes

$$E_2 = E_1$$

Where $E = E_K + E_P$ is defined as the total energy.

SAMPLE 12.8 Basic calculations: Find $\vec{L}, \dot{\vec{L}}, \vec{H}_{/C}, \dot{\vec{H}}_{/C}, E_K, \dot{E}_K$ for a given particle P with mass $m_P = 1$ kg, given position, velocity, acceleration, and a point C. Specifically, we are given $\vec{r}_P = (\hat{i} + \hat{j} + \hat{k})$ m, $\vec{v}_P = 3$ m/s $(\hat{i} + \hat{j})$, $\vec{a}_P = 2$ m/s² $(\hat{i} - \hat{j} - \hat{k})$, and $\vec{r}_C = (2\hat{i} + \hat{k})$ m.

Solution Since $\vec{r}_P = (\hat{i} + \hat{j} + \hat{k})$ m and $\vec{r}_C = (2\hat{i} + \hat{k})$ m,

$$\vec{r}_{P/C} = \vec{r}_P - \vec{r}_C = (-\hat{i} + \hat{j}) \text{ m}.$$

So we have the motion quantities

$$\begin{aligned}\vec{L} &= m\vec{v}_P \\ &= (1 \text{ kg}) \cdot [(3 \text{ m/s})(\hat{i} + \hat{j})] \\ &= 3(\hat{i} + \hat{j}) \frac{\text{kg} \cdot \text{m}}{\text{s}} \\ &= 3 \text{ N} \cdot \text{s}(\hat{i} + \hat{j})\end{aligned}$$

$$\begin{aligned}\dot{\vec{L}} &= m\vec{a}_P \\ &= (1 \text{ kg})[(2 \text{ m/s}^2)(\hat{i} - \hat{j} - \hat{k})] \\ &= 2(\hat{i} - \hat{j} - \hat{k}) \frac{\text{kg} \cdot \text{m}}{\text{s}^2} \\ &= 2 \text{ N}(\hat{i} - \hat{j} - \hat{k})\end{aligned}$$

$$\begin{aligned}\vec{H}_{/C} &= \vec{r}_{P/C} \times m\vec{v}_P \\ &= [(-\hat{i} + \hat{j}) \text{ m}] \times [(1 \text{ kg})3 \text{ m/s}(\hat{i} + \hat{j})] \\ &= -(6 \text{ kg} \cdot \text{m}^2/\text{s})\hat{k}\end{aligned}\tag{12.11}$$

$$\begin{aligned}\dot{\vec{H}}_{/C} &= \vec{r}_{P/C} \times m\vec{a}_P \\ &= [(-\hat{i} + \hat{j}) \text{ m}] \times [(1 \text{ kg})2 \text{ m/s}^2(\hat{i} - \hat{j} - \hat{k})] \\ &= -(2 \text{ kg} \cdot \text{m}^2/\text{s}^2)(\hat{i} + \hat{j})\end{aligned}$$

$$\begin{aligned}E_K &= \frac{1}{2}m|\vec{v}_P|^2 \\ &= \frac{1}{2}(1 \text{ kg})(3\sqrt{2} \text{ m/s})^2 \\ &= 9 \frac{\text{kg} \cdot \text{m}^2}{\text{s}^2} \\ &= 9 \text{ N} \cdot \text{m}\end{aligned}$$

$$\begin{aligned}\dot{E}_K &= \frac{d}{dt}\left(\frac{m}{2}\vec{v}_P \cdot \vec{v}_P\right) \\ &= \frac{m}{2}[\vec{v}_P \cdot \dot{\vec{v}}_P + \dot{\vec{v}}_P \cdot \vec{v}_P] \\ &= m\vec{v}_P \cdot \vec{a}_P \\ &= 1 \text{ kg}[(3 \text{ m/s})(\hat{i} + \hat{j})] \cdot [(2 \text{ m/s}^2)(\hat{i} - \hat{j} - \hat{k})] \\ &= 0.\end{aligned}$$

Note: $\frac{d}{dt}\left(\frac{1}{2}v^2\right) \neq |\dot{\vec{v}}| |\vec{a}|$.




Figure 12.26

SAMPLE 12.9 Direct application of the formulas: A 2 kg block is moving with a velocity $\vec{v}(t) = u_0 e^{-ct} \hat{i} + v_0 \hat{j}$, where $u_0 = 5 \text{ m/s}$, $v_0 = 10 \text{ m/s}$, and $c = 0.5/\text{s}$. Consider the time interval between $t_1 = 1 \text{ s}$ to $t_2 = 3 \text{ s}$.

1. Find the net change in the linear momentum of the block, $\Delta \vec{L} = \vec{L}(t_2) - \vec{L}(t_1)$.
2. Find the force $\vec{F}(t)$ on the block and compute the impulse $\int_{t_1}^{t_2} \vec{F} dt$ and show that it is the same as $\Delta \vec{L}$ computed above.
3. Find the change in kinetic energy from direct computation of energy and compare with work done by computing $\int_{t_1}^{t_2} P dt$.

Solution

1. For the given block we have, $\vec{L} = m\vec{v} = m(u_0 e^{-ct} \hat{i} + v_0 \hat{j})$. Therefore,

$$\Delta L = \vec{L}(t_2) - \vec{L}(t_1) = m u_0 (e^{-ct_2} - e^{-ct_1}) \hat{i}.$$

Substituting the given values, $m = 2 \text{ kg}$, $u_0 = 5 \text{ m/s}$, $v_0 = 10 \text{ m/s}$, $c = 0.5/\text{s}$, $t_1 = 1 \text{ s}$, and $t_2 = 3 \text{ s}$ we get

$$\Delta \vec{L} = 2 \text{ kg} \cdot 5 \text{ m/s} (e^{-0.5/\text{s} \cdot 3 \text{ s}} - e^{-0.5/\text{s} \cdot 1 \text{ s}}) \hat{i} = -(3.83 \text{ kg} \cdot \text{m/s}) \hat{i}.$$

$$\Delta \vec{L} = -(3.83 \text{ kg} \cdot \text{m/s}) \hat{i}$$

2. To calculate the impulse, $\int \vec{F} dt$, we need to find the force first. Since $\vec{F} = m\vec{a} = m\dot{\vec{v}}$, we get

$$\vec{F}(t) = m \frac{d}{dt} (u_0 e^{-ct} \hat{i} + v_0 \hat{j}) = -m c u_0 e^{-ct} \hat{i}.$$

Hence, the impulse is

$$\begin{aligned} \int_{t_1}^{t_2} \vec{F} dt &= - \int_{t_1}^{t_2} m c u_0 e^{-ct} dt \hat{i} = m u_0 (e^{-ct_2} - e^{-ct_1}) \hat{i} \\ &= 2 \text{ kg} \cdot 5 \text{ m/s} (e^{-0.5/\text{s} \cdot 3 \text{ s}} - e^{-0.5/\text{s} \cdot 1 \text{ s}}) \hat{i} \\ &= -3.83 \text{ kg} \cdot \text{m/s} \hat{i} \end{aligned}$$

which is, expectedly, the same answer as obtained above for ΔL .

3. To find the kinetic energy, we need the speed of the particle, $v = |\vec{v}| = \sqrt{v_x^2 + v_y^2}$. Now, the change in kinetic energy is

$$\begin{aligned} \Delta E_K &= E_{K2} - E_{K1} = \frac{1}{2} m (v_2^2 - v_1^2) \\ &= \frac{1}{2} m (u_0^2 e^{-2ct_2} + v_0^2 - u_0^2 e^{-2ct_1} - v_0^2) \\ &= \frac{1}{2} m u_0^2 (e^{-2ct_2} - e^{-2ct_1}) = -7.95 \text{ N} \cdot \text{m}. \end{aligned}$$

Now, we can compare this value by computing the work done $\int P dt$, since $\Delta E_K = \int P dt$. To compute the power $P = \vec{F} \cdot \vec{v}$, we need to find the dot product between the force and the velocity. Since $\vec{F} = -m c u_0 e^{-ct} \hat{i}$, and $\vec{v} = u_0 e^{-ct} \hat{i} + v_0 \hat{j}$, we get, $\vec{F} \cdot \vec{v} = -m c u_0^2 e^{-2ct}$. Therefore, the work done is,

$$\begin{aligned} W &= \int_{t_1}^{t_2} P dt = \int_{t_1}^{t_2} -m c u_0^2 e^{-2ct} dt \\ &= -m c u_0^2 (e^{-2ct_2} - e^{-2ct_1}) = -7.95 \text{ N} \cdot \text{m}. \end{aligned}$$

$$\Delta E_K = -7.95 \text{ N} \cdot \text{m}, \quad W = -7.95 \text{ N} \cdot \text{m}$$




Figure 12.27: The plot of speed $v = |\vec{v}| = \sqrt{v_x^2 + v_y^2}$ vs time. The speeds at t_1 and t_2 are of interest for computing the kinetic energy at the two instants.




Figure 12.28: The area under the $P(t)$ curve between t_1 and t_2 is the work done $W = \int_{t_1}^{t_2} P dt$.

SAMPLE 12.10 Angular momentum: direct application of the formula. The position of a particle of mass $m = 0.5 \text{ kg}$ is $\vec{r}(t) = \ell \sin(\lambda t)\hat{i} + h\hat{j}$; where $\lambda = \pi/2 \text{ rad/s}$, $h = 2 \text{ m}$, $\ell = 2 \text{ m}$, and $\vec{r}$ is measured from the origin.

1. Find the net change in linear momentum $\Delta \vec{L}$ of the particle between $t = 1 \text{ s}$ and $t = 3 \text{ s}$.
2. Find the net change in angular momentum $\Delta \vec{H}_{/O}$ of the particle about the origin between $t = 1 \text{ s}$ and $t = 3 \text{ s}$.
3. Find the angular impulse $\int_{1\text{s}}^{3\text{s}} M dt$ about the origin and compare the result with $\Delta \vec{H}_{/O}$ found above.

Solution

1. Linear momentum: Let the two instants of interest be $t_1 (= 1 \text{ s})$ and $t_2 (= 3 \text{ s})$. The net change in linear momentum, $\Delta \vec{L} = \vec{L}_2 - \vec{L}_1 = m(\vec{v}_2 - \vec{v}_1)$. Since $\vec{v} = \dot{\vec{r}} = \ell \lambda \cos(\lambda t)\hat{i}$, we get

$$\begin{aligned}\Delta \vec{L} &= m(\vec{v}_2 - \vec{v}_1) = m \ell \lambda (\cos \lambda t_2 - \cos \lambda t_1)\hat{i} \\ &= (0.5 \text{ kg})(2 \text{ m})\left(\frac{\pi}{2} \text{ rad/s}\right)\left(\cos \frac{\pi}{2} - \cos \frac{3\pi}{2}\right)\hat{i} \\ &= \vec{0}.\end{aligned}$$

The answer makes sense because both $\vec{v}_1 = \vec{0}$ and $\vec{v}_2 = \vec{0}$. In fact, finding the velocity at $t_1 = 1 \text{ s}$ and $t_2 = 3 \text{ s}$ would have made the calculation much simpler.

$$\Delta \vec{L} = \vec{0}$$

2. Angular momentum: The net change in angular momentum between t_1 and t_2 is,

$$\Delta \vec{H}_{/O} = (\vec{H}_{/O})_2 - (\vec{H}_{/O})_1 = \vec{r}_{/O} \times m \underbrace{\vec{v}_2}_{\vec{0}} - \vec{r}_{/O} \times m \underbrace{\vec{v}_1}_{\vec{0}} = \vec{0}.$$

$$\Delta \vec{H}_{/O} = \vec{0}$$

Note that it so happens that velocities at the two instants are zero and hence, both $(\vec{H}_{/O})_1$ and $(\vec{H}_{/O})_2$ are zero, making $\Delta \vec{H}_{/O}$ also zero. It is, however, possible that we could get $\Delta \vec{H}_{/O}$ to be zero even if the $(\vec{H}_{/O})_1$ and $(\vec{H}_{/O})_2$ were non-zero (when they are equal).

3. Moment impulse: Now, let us find the impulse due to the moment, $\int M dt$ between the two given time instants and see if that matches with the net zero change in angular momentum. We first need to compute the moment $\vec{M}_O = \vec{r}_{/O} \times \vec{F} = \vec{r}_{/O} \times m\vec{a}$:

$$\vec{M}_{/O}(t) = \vec{r}_{/O} \times m\vec{a} = (\ell \sin(\lambda t)\hat{i} + h\hat{j}) \times m(-\ell \lambda^2 \sin \lambda t\hat{i}) = m \ell h \lambda^2 \sin \lambda t \hat{k}.$$

Therefore, the impulse due to this moment is

$$\begin{aligned}\int_{t_1}^{t_2} \vec{M}_{/O} dt &= \int_{1\text{s}}^{3\text{s}} (m \ell h \lambda^2 \sin \lambda t \hat{k}) dt = m \ell h \lambda^2 \hat{k} \int_{1\text{s}}^{3\text{s}} \sin(\lambda t) dt \\ &= m \ell h \lambda^2 \hat{k} \left[-\frac{\cos \lambda t}{\lambda} \right]_{1\text{s}}^{3\text{s}} = m \ell h \lambda \hat{k} \left[\cos \frac{3\pi}{2} - \cos \frac{\pi}{2} \right] \\ &= \vec{0}\end{aligned}$$

as expected. It can also be seen from a plot of $|\vec{M}_{/O}|$ vs t , as shown in fig. 12.30, that the net area under the moment between t_1 and t_2 is zero, giving a zero moment impulse.




Figure 12.29




Figure 12.30: Plot of $M(t)$ where $\vec{M}_{/O}(t) = M(t)\hat{k}$. The area under the moment curve between t_1 and t_2 is the magnitude of the moment impulse.

12.2 Momentum and energy for particle motion

Preparatory Problems

12.2.1 What symbols do we use for the following quantities? What are the definitions of these quantities? Which are vectors and which are scalars? What are the SI and US standard units for the following quantities?

- linear momentum
- rate of change of linear momentum
- angular momentum
- rate of change of angular momentum
- kinetic energy
- rate of change of kinetic energy
- moment
- work
- power

12.2.2 Does angular momentum depend on reference point? (Assume that all candidate points are fixed in the same Newtonian reference frame.)

12.2.3 Does kinetic energy depend on reference point? (Assume that all candidate points are fixed in the same Newtonian reference frame.)

12.2.4 What is the relation between the dynamics 'Linear Momentum Balance' equation and the statics 'Force Balance' equation?

12.2.5 What is the relation between the dynamics 'Angular Momentum Balance' equation and the statics 'Moment Balance' equation?

12.2.6 A ball of mass $m = 0.1$ kg is thrown from a height of $h = 10$ m above the ground with velocity $\vec{v} = 120 \text{ km/h} \hat{i} - 120 \text{ km/h} \hat{j}$. What is the kinetic energy of the ball at its release?

12.2.7 A ball of mass $m = 0.2$ kg is thrown from a height of $h = 20$ m above the ground with velocity $\vec{v} = 120 \text{ km/h} \hat{i} - 120 \text{ km/h} \hat{j} - 10 \text{ km/h} \hat{k}$. What is the kinetic energy of the ball at its release?

12.2.8 How do you calculate P , the power of all external forces acting on a particle, from the forces $\vec{F}_i$ and the velocity $\vec{v}$ of the particle?

12.2.9 A particle A has velocity $\vec{v}_A$ and mass m_A . A particle B has velocity $\vec{v}_B = 2 \vec{v}_A$ and mass equal to the other $m_B = m_A$. What is the relationship between:

- $\vec{L}_A$ and $\vec{L}_B$,
- $\vec{H}_{A/C}$ and $\vec{H}_{B/C}$, and
- E_{KA} and E_{KB} ?

12.2.10 A bullet of mass 50 g travels with a velocity $\vec{v} = 0.8 \text{ km/s} \hat{i} + 0.6 \text{ km/s} \hat{j}$. (a) What is the linear momentum of the bullet? (Answer in consistent units.)

12.2.11 A particle has position $\vec{r} = 4 \text{ m} \hat{i} + 7 \text{ m} \hat{j}$, velocity $\vec{v} = 6 \text{ m/s} \hat{i} - 3 \text{ m/s} \hat{j}$, and acceleration $\vec{a} = -2 \text{ m/s}^2 \hat{i} + 9 \text{ m/s}^2 \hat{j}$. For each position of a point P defined below, find $\vec{H}_P$, the angular momentum of the particle with respect to the point P .

- $\vec{r}_P = 4 \text{ m} \hat{i} + 7 \text{ m} \hat{j}$,
- $\vec{r}_P = -2 \text{ m} \hat{i} + 7 \text{ m} \hat{j}$, and
- $\vec{r}_P = 0 \text{ m} \hat{i} + 7 \text{ m} \hat{j}$,
- $\vec{r}_P = \vec{0}$

12.2.12 The position vector of a particle of mass 1 kg at an instant t is $\vec{r} = 2 \text{ m} \hat{i} - 0.5 \text{ m} \hat{j}$. If the velocity of the particle at this instant is $\vec{v} = -4 \text{ m/s} \hat{i} + 3 \text{ m/s} \hat{j}$, compute (a) the linear momentum $\vec{L} = m\vec{v}$ and (b) the angular momentum ($\vec{H}_O = \vec{r}_O \times (m\vec{v})$).

12.2.13 The position of a particle of mass $m = 0.5$ kg is $\vec{r}(t) = \ell \sin(\omega t) \hat{i} + h \hat{j}$; where $\omega = 2 \text{ rad/s}$, $h = 2 \text{ m}$, $\ell = 2 \text{ m}$, and $\vec{r}$ is measured from the origin.


- Find the kinetic energy of the particle at $t = 0$ s and $t = 5$ s.
- Find the rate of change of kinetic energy at $t = 0$ s and $t = 5$ s.

12.2.14 For a particle

$$E_K = \frac{1}{2} m v^2.$$

Why does it follow that $\dot{E}_K = m \vec{v} \cdot \vec{a}$? [hint: write v^2 as $\vec{v} \cdot \vec{v}$ and then use the product rule of differentiation.]

12.2.15 Consider a projectile of mass m at some instant in time t during its flight. Let $\vec{v}$ be the velocity of the projectile at this instant (see the figure). In addition to the force of gravity, a drag force acts on the projectile. The drag force is proportional to the square of the speed (speed $\equiv |\vec{v}| = v$) and acts in the opposite direction. Find an expression for the net power of these forces ($P = \sum \vec{F} \cdot \vec{v}$) on the particle.



Problem 12.2.15

12.2.16 A 10 gm wad of paper is tossed into the air. At some instant, the position, velocity, and acceleration of its center of mass are $\vec{r} = 3 \text{ m} \hat{i} + 3 \text{ m} \hat{j} + 6 \text{ m} \hat{k}$, $\vec{v} = -9 \text{ m/s} \hat{i} + 24 \text{ m/s} \hat{j} + 30 \text{ m/s} \hat{k}$, and $\vec{a} = -10 \text{ m/s}^2 \hat{i} + 24 \text{ m/s}^2 \hat{j} + 32 \text{ m/s}^2 \hat{k}$, respectively. What is the translational kinetic energy of the wad at the instant of interest?

12.2.17 A 2 kg particle moves so that its position $\vec{r}$ is given by

$$\vec{r}(t) = [5 \sin(at) \hat{i} + bt^2 \hat{j} + ct \hat{k}] \text{ meters}$$

where $a = \pi / \text{sec}$, $b = .25 / \text{sec}^2$, $c = 2 / \text{sec}$.

- What is the linear momentum of the particle at $t = 1$ sec?

- b) What is the force acting on the particle at $t = 1$ sec?

12.2.18 A particle A has mass m_A and velocity $\vec{v}_A$. A particle B at the same location has mass $m_B = 2m_A$ and velocity equal to the other $\vec{v}_B = \vec{v}_A$. Point C is a reference point. What is the relationship between:

- $\vec{L}_A$ and $\vec{L}_B$,
- $\vec{H}_{A/C}$ and $\vec{H}_{B/C}$, and
- E_{KA} and E_{KB} ?

12.2.19 A particle of mass $m = 3$ kg moves in space. Its position, velocity, and acceleration at a given time are $\vec{r} = 2m\hat{i} + 3m\hat{j} + 5m\hat{k}$, $\vec{v} = -3m/s\hat{i} + 8m/s\hat{j} + 10m/s\hat{k}$, and $\vec{a} = -5m/s^2\hat{i} + 12m/s^2\hat{j} + 16m/s^2\hat{k}$, respectively. For this particle at the instant of interest, find its:

- linear momentum $\vec{L}$,
- rate of change of linear momentum $\dot{\vec{L}}$,
- angular momentum about the origin $\vec{H}_O$,
- rate of change of angular momentum about the origin $\dot{\vec{H}}_O$,
- kinetic energy E_K , and
- rate of change of kinetic energy $\dot{E}_K$.

12.2.20 A particle has position $\vec{r} = 3m\hat{i} - 2m\hat{j} + 4m\hat{k}$, velocity $\vec{v} = 2m/s\hat{i} - 3m/s\hat{j} + 7m/s\hat{k}$, and acceleration $\vec{a} = 1m/s^2\hat{i} - 8m/s^2\hat{j} + 3m/s^2\hat{k}$. For each position of a point P defined below, find the rate of change of angular momentum, $\dot{\vec{H}}_P$, of the particle with respect to the point P .


- $\vec{r}_P = 3m\hat{i} - 2m\hat{j} + 4m\hat{k}$,
- $\vec{r}_P = 6m\hat{i} - 4m\hat{j} + 8m\hat{k}$,
- $\vec{r}_P = -9m\hat{i} + 6m\hat{j} - 12m\hat{k}$, and
- $\vec{r}_P = \vec{0}$

More-Involved Problems

12.2.21 A particle of mass $m = 6$ kg is moving in space. Its position, velocity, and acceleration at some instant are $\vec{r} = 1m\hat{i} - 2m\hat{j} + 4m\hat{k}$, $\vec{v} = 3m/s\hat{i} + 4m/s\hat{j} - 7m/s\hat{k}$, and $\vec{a} = 5m/s^2\hat{i} + 11m/s^2\hat{j} - 9m/s^2\hat{k}$, respectively. At this instant, find:

- the net force $\sum \vec{F}$ on the particle,
- the net moment on the particle about the origin $\sum \vec{M}_O$ due to the applied forces, and
- the power P of the applied forces.

Particle FBD



Problem 12.2.21: FBD of the particle

12.2.22 At a time of interest, a particle with mass $m_1 = 5$ kg has position, velocity, and acceleration $\vec{r}_1 = 3m\hat{i}$, $\vec{v}_1 = -4m/s\hat{j}$, and $\vec{a}_1 = 6m/s^2\hat{j}$, respectively. Another particle with mass $m_2 = 5$ kg has position, velocity, and acceleration $\vec{r}_2 = -6m\hat{i}$, $\vec{v}_2 = 5m/s\hat{j}$, and $\vec{a}_2 = -4m/s^2\hat{j}$, respectively. For this system of two particles, and at this time, find its

- linear momentum $\vec{L}$,
- rate of change of linear momentum $\dot{\vec{L}}$
- angular momentum about the origin $\vec{H}_O$,
- rate of change of angular momentum about the origin $\dot{\vec{H}}_O$,
- kinetic energy E_K , and
- rate of change of kinetic energy $\dot{E}_K$.

12.2.23 A particle of mass $m = 250$ gm is shot straight up (parallel to the y -axis) from the x -axis at a distance $d = 2$ m from the origin. The velocity of the particle is given by $\vec{v} = v\hat{j}$ where $v^2 = v_0^2 - 2ah$, $v_0 = 100$ m/s, $a = 10$ m/s² and h is the height of the particle from the x -axis.

- Find the linear momentum of the particle at the outset of motion ($h = 0$).
- Find the angular momentum of the particle about the origin at the outset of motion ($h = 0$).
- Find the linear momentum of the particle when the particle is 20 m above the x -axis.
- Find the angular momentum of the particle about the origin when the particle is 20 m above the x -axis.

12.2.2 Does angular momentum depend on reference point? (Assume that all candidate points are fixed in the same Newtonian reference frame.)

11.2.2

No,

$\vec{L} = \frac{d}{dt}(m\vec{r}\vec{v})$, is the same with respect to any point in the same Newtonian reference frame in which $\vec{v}$ is measured.

12.2.4 What is the relation between the dynamics 'Linear Momentum Balance' equation and the statics 'Force Balance' equation?

11.2.4

$$\vec{F}_{\text{net}} = \sum \vec{F} = \frac{d(\vec{L})}{dt} = \dot{\vec{L}}$$

Disclaimer: We have lost track of the many people who wrote these solutions. They are not official and have not been reviewed by the authors for correctness.

12.2.5 What is the relation between the dynamics 'Angular Momentum Balance' equation and the statics 'Moment Balance' equation?

12.2.5

$$\vec{M}_{\text{net}/c} = \frac{d}{dt} (\vec{H}_{/c}) = \dot{\vec{H}}_{/c}$$

12.2.6 A ball of mass $m = 0.1 \text{ kg}$ is thrown from a height of $h = 10 \text{ m}$ above the ground with velocity $\vec{v} = 120 \text{ km/h} \hat{i} - 120 \text{ km/h} \hat{j}$. What is the kinetic energy of the ball at its release?

12.2.6

$$m = 0.1 \text{ kg};$$

$$h = 10 \text{ m}$$

$$\vec{V} = 120 \frac{\text{km}}{\text{hr}} \hat{i} - 120 \frac{\text{km}}{\text{hr}} \hat{j}$$

$$\vec{V} = \frac{100}{3} [\hat{i} - \hat{j}] \text{ m/sec}$$

$$KE = \frac{1}{2} m \vec{V} \cdot \vec{V} = \frac{0.1}{2} \left[\left(\frac{100}{3} \right)^2 + \left(\frac{100}{3} \right)^2 \right] \text{ Joules}$$

$$KE = 111.11 \text{ Joules}$$

Disclaimer: We have lost track of the many people who wrote these solutions. They are not official and have not been reviewed by the authors for correctness.

12.2.7 A ball of mass $m = 0.2 \text{ kg}$ is thrown from a height of $h = 20 \text{ m}$ above the ground with velocity $\vec{v} = 120 \text{ km/h} \hat{i} - 120 \text{ km/h} \hat{j} - 10 \text{ km/h} \hat{k}$. What is the kinetic energy of the ball at its release?

12.2.7

$$m = 0.2 \text{ kg}, \quad h = 20 \text{ m}$$

$$\vec{v} = (120 \hat{i} - 120 \hat{j} - 10 \hat{k}) \text{ km/hr}$$

$$KE = \frac{1}{2} m \vec{v} \cdot \vec{v} = 0.1 [v_x^2 + v_y^2 + v_z^2]$$

$$= 0.1 \left[\left(\frac{1000}{3}\right)^2 + \left(\frac{1000}{3}\right)^2 + (2.77)^2 \right]$$

$$KE = 222.989 \text{ Joules.}$$

12.2.8 How do you calculate P , the power of all external forces acting on a particle, from the forces $\vec{F}_i$ and the velocity $\vec{v}$ of the particle?

11.2.8

$$\text{Power} = \sum_{i=1}^{n_1} \frac{d}{dt} (\vec{F}_i \cdot d\vec{r}) = \sum_{i=1}^{n_1} \vec{F}_i \cdot \vec{v}$$

Disclaimer: We have lost track of the many people who wrote these solutions. They are not official and have not been reviewed by the authors for correctness.

12.2.9 A particle A has velocity $\vec{v}_A$ and mass m_A . A particle B has velocity $\vec{v}_B = 2\vec{v}_A$ and mass equal to the other $m_B = m_A$. What is the relationship between:

- $\vec{L}_A$ and $\vec{L}_B$,
- $\vec{H}_{A/C}$ and $\vec{H}_{B/C}$, and
- E_{KA} and E_{KB} ?

$$m_B = m_A, \quad \vec{v}_B = 2\vec{v}_A$$

$$a) \vec{L}_B = m_B \vec{v}_B = 2m_A \vec{v}_A = 2\vec{L}_A$$

$$\vec{L}_B = 2\vec{L}_A$$

$$b) \vec{H}_{B/C} = \vec{r}_{B/C} \times \vec{L}_B = 2\vec{r}_{B/C} \times \vec{L}_A$$

$$\vec{H}_{A/C} = \vec{r}_{A/C} \times \vec{L}_A$$

$$\text{If } \vec{r}_{B/C} = \vec{r}_{A/C}$$

then,

$$\vec{H}_{B/C} = 2\vec{H}_{A/C}$$

$$c) E_{KA} = \frac{1}{2} m_A \vec{v}_A \cdot \vec{v}_A, \quad E_{KB} = \frac{1}{2} m_B \vec{v}_B \cdot \vec{v}_B$$

$$E_{KB} = \frac{1}{2} m_B \vec{v}_B \cdot \vec{v}_B = \frac{1}{2} m_A (2\vec{v}_A) \cdot (2\vec{v}_A) = 4E_{KA}$$

$$E_{KB} = 4E_{KA}$$

12.2.10 A bullet of mass 50 g travels with a velocity $\vec{v} = 0.8 \text{ km/s} \hat{i} + 0.6 \text{ km/s} \hat{j}$. (a) What is the linear momentum of the bullet? (Answer in consistent units.)

$$m = 50 \text{ grams}$$

$$\vec{v} = [0.8 \hat{i} + 0.6 \hat{j}] \text{ km/sec}$$

Linear momentum of the Bullet

$$\vec{L} = m\vec{v} = \frac{50}{1000} [0.8 \hat{i} + 0.6 \hat{j}] \text{ km/sec}$$

$$\vec{L} = [40 \hat{i} + 30 \hat{j}] \text{ kg m/sec}$$

12.2.11 A particle has position $\vec{r} = 4\text{ m}\hat{i} + 7\text{ m}\hat{j}$, velocity $\vec{v} = 6\text{ m/s}\hat{i} - 3\text{ m/s}\hat{j}$, and acceleration $\vec{a} = -2\text{ m/s}^2\hat{i} + 9\text{ m/s}^2\hat{j}$. For each position of a point P defined below, find $\vec{H}_P$, the angular momentum of the particle with respect to

the point P.

- $\vec{r}_P = 4\text{ m}\hat{i} + 7\text{ m}\hat{j}$,
- $\vec{r}_P = -2\text{ m}\hat{i} + 7\text{ m}\hat{j}$, and
- $\vec{r}_P = 0\text{ m}\hat{i} + 7\text{ m}\hat{j}$,
- $\vec{r}_P = \vec{0}$

32

11.2.11

$$\vec{r} = (4\hat{i} + 7\hat{j})\text{ m}$$

$$\vec{v} = (6\hat{i} - 3\hat{j})\text{ m/sec}$$

$$\vec{a} = (-2\hat{i} + 9\hat{j})\text{ m/sec}^2$$

$$\text{say mass} = m$$

$$\vec{H}_P = \vec{r}_{P/P} \times \vec{L} = m \vec{r}_{P/P} \times \vec{v}$$

$$\text{a) } \vec{r}_P = (4\hat{i} + 7\hat{j})\text{ m}$$

$$\begin{aligned} \vec{H}_P &= m(4\hat{i} + 7\hat{j}) \times (6\hat{i} - 3\hat{j})\text{ kg m}^2/\text{sec} \\ &= m(-12 - 42)\hat{k}\text{ kg m}^2/\text{sec} \end{aligned}$$

$$\vec{H}_P = -54m\hat{k}\text{ kg m}^2/\text{sec}$$

$$\text{b) } \vec{r}_P = (-2\hat{i} + 7\hat{j})\text{ m}$$

$$\begin{aligned} \vec{H}_P &= m(-2\hat{i} + 7\hat{j}) \times (6\hat{i} - 3\hat{j})\text{ kg m}^2/\text{sec} \\ &= m(6 - 42)\hat{k}\text{ kg m}^2/\text{sec} \end{aligned}$$

$$\vec{H}_P = -36m\hat{k}\text{ kg m}^2/\text{sec}$$

$$\text{c) } \vec{H}_P = m(7\hat{j}) \times (6\hat{i} - 3\hat{j})\text{ kg m}^2/\text{sec}$$

$$\vec{H}_P = -42m\hat{k}\text{ kg m}^2/\text{sec}$$

$$\text{d) } \vec{H}_P = m(\vec{0}) \times (6\hat{i} - 3\hat{j})$$

$$\vec{H}_P = 0\text{ kg m}^2/\text{sec}$$

Disclaimer: We have lost track of the many people who wrote these solutions. They are not official and have not been reviewed by the authors for correctness.

12.2.12 The position vector of a particle of mass 1 kg at an instant t is $\vec{r} = 2\text{ m}\hat{i} - 0.5\text{ m}\hat{j}$. If the velocity of the particle at this instant is $\vec{v} = -4\text{ m/s}\hat{i} + 3\text{ m/s}\hat{j}$, compute (a) the linear momentum $\vec{L} = m\vec{v}$ and (b) the angular momentum ($\vec{H}_{/O} = \vec{r}_O \times (m\vec{v})$).

(33)

11.2.12

$$m = 1 \text{ kg}, \quad \vec{r} = (2\hat{i} - 0.5\hat{j})$$

$$\vec{v} = -4\hat{i} + 3\hat{j}$$

a) Linear Momentum,

$$\vec{L} = m\vec{v} = (-4\hat{i} + 3\hat{j}) \text{ kg m/sec}$$

b) $\vec{H}_{/O} = m\vec{r} \times \vec{v}$

$$= (2\hat{i} - 0.5\hat{j}) \times (-4\hat{i} + 3\hat{j}) \text{ kg m}^2/\text{sec}^2$$

$$= (6 - 2)\hat{k} \text{ kg m}^2/\text{sec}$$

$$\vec{H}_{/O} = 4\hat{k} \text{ kg m}^2/\text{sec}$$

12.2.13 The position of a particle of mass $m = 0.5 \text{ kg}$ is $\vec{r}(t) = \ell \sin(\omega t) \hat{i} + h \hat{j}$; where $\omega = 2 \text{ rad/s}$, $h = 2 \text{ m}$, $\ell = 2 \text{ m}$, and $\vec{r}$ is measured from the origin.

- a) Find the kinetic energy of the particle at $t = 0 \text{ s}$ and $t = 5 \text{ s}$.
 b) Find the rate of change of kinetic energy at $t = 0 \text{ s}$ and $t = 5 \text{ s}$.

(34)

11.2.13

$$m = 0.5 \text{ kg}$$

$$\vec{r}(t) = \ell \sin \omega t \hat{i} + h \hat{j}$$

$$\vec{v}(t) = \frac{d\vec{r}}{dt} = \omega \ell \cos \omega t \hat{i}$$

$$\omega = 2 \text{ rad/sec}$$

$$h = 2 \text{ m}$$

$$\ell = 2 \text{ m}$$

$$a) E_K = \frac{1}{2} m \vec{v} \cdot \vec{v}$$

$$= \frac{1}{2} m (\omega \ell \cos \omega t)^2 = \frac{1}{2} m \omega^2 \ell^2 \cos^2 \omega t = 4 \cos^2 \omega t$$

$$\text{at } t = 0$$

$$E_K = 4 \text{ Joules}$$

$$\text{at } t = 5 \text{ sec}$$

$$E_K = 4 [\cos(2 \times 5)]^2$$

$$E_K = 2.8161 \text{ Joules}$$

$$b) \dot{E}_K = \frac{d}{dt} \left[\frac{1}{2} m \omega^2 \ell^2 \cos^2 \omega t \right]$$

$$= 4 \frac{d}{dt} \cos^2 \omega t = -8 \cos \omega t \sin \omega t = -4 \sin 2 \omega t$$

$$\text{at } t = 0$$

$$\dot{E}_K = 0 \text{ Joules/sec}$$

$$\text{at } t = 5 \text{ sec}$$

$$\dot{E}_K = -4 \sin(20)$$

$$\dot{E}_K = -3.6517 \text{ Joules/sec}$$

Disclaimer: We have lost track of the many people who wrote these solutions. They are not official and have not been reviewed by the authors for correctness.

12.2.14 For a particle


$$E_K = \frac{1}{2} m v^2.$$

Why does it follow that $\dot{E}_K = m \vec{v} \cdot \vec{a}$?
[hint: write v^2 as $\vec{v} \cdot \vec{v}$ and then use the product rule of differentiation.]

$$\begin{aligned} E_K &= \frac{1}{2} m \vec{v} \cdot \vec{v} \\ \dot{E}_K &= \frac{d}{dt} \left(\frac{1}{2} m \vec{v} \cdot \vec{v} \right) \\ &= \frac{1}{2} m \left[\vec{v} \cdot \frac{d\vec{v}}{dt} + \frac{d\vec{v}}{dt} \cdot \vec{v} \right] \\ &= m \left[\vec{v} \cdot \frac{d\vec{v}}{dt} \right] \\ \dot{E}_K &= m \vec{v} \cdot \vec{a} \end{aligned}$$


12.2.15 Consider a projectile of mass m at some instant in time t during its flight. Let $\vec{v}$ be the velocity of the projectile at this instant (see the figure). In addition to the force of gravity, a drag force acts on the projectile. The drag force is proportional to the square of the speed (speed $\equiv |\vec{v}| = v$) and acts in the opposite direction. Find an expression for the

net power of these forces ($P = \sum \vec{F} \cdot \vec{v}$) on the particle.



Problem 12.15

11.2.15



Say $\vec{v} = v_x \hat{i} + v_y \hat{j}$

$$F_D = -bv^2 \hat{e}_r = -bv^2 \frac{\vec{v}}{v} = -bv\vec{v}, \quad v = \sqrt{v_x^2 + v_y^2}$$

Net Power of force

$$P = \sum \vec{F} \cdot \vec{v}$$

$$= -(bv) \vec{v} \cdot \vec{v} - mg \hat{j} \cdot \vec{v}$$

$$= -bv^3 - mgv_y$$

$$= -bv^3 - mg(v \sin \theta)$$

$$P = -mgv \sin \theta - bv^3$$

12.2.16 A 10 gm wad of paper is tossed into the air. At some instant, the position, velocity, and acceleration of its center of mass are $\vec{r} = 3 \text{ m}\hat{i} + 3 \text{ m}\hat{j} + 6 \text{ m}\hat{k}$,

$\vec{v} = -9 \text{ m/s}\hat{i} + 24 \text{ m/s}\hat{j} + 30 \text{ m/s}\hat{k}$, and $\vec{a} = -10 \text{ m/s}^2\hat{i} + 24 \text{ m/s}^2\hat{j} + 32 \text{ m/s}^2\hat{k}$, respectively. What is the translational kinetic energy of the wad at the instant of interest?

$$m = 10 \text{ gms}$$

$$E_k = \frac{1}{2} m \vec{v} \cdot \vec{v} = \frac{1}{2} \left(\frac{10}{1000} \right) (9^2 + 24^2 + 30^2) \text{ Joules}$$

$$E_k = 7.785 \text{ Joules}$$

12.2.17 A 2 kg particle moves so that its position $\vec{r}$ is given by

$$\vec{r}(t) = [5 \sin(at)\hat{i} + bt^2\hat{j} + ct\hat{k}] \text{ meters}$$

where $a = \pi/\text{sec}$, $b = .25/\text{sec}^2$, $c = 2/\text{sec}$.

a) What is the linear momentum of the particle at $t = 1 \text{ sec}$?

b) What is the force acting on the particle at $t = 1 \text{ sec}$?

$$\begin{aligned}
 m &= 2 \text{ kg} \\
 \vec{r} &= [5 \sin at \hat{i} + bt^2 \hat{j} + ct \hat{k}] \\
 \vec{v} &= 5a \cos at \hat{i} + 2bt \hat{j} + c \hat{k} \\
 \vec{a} &= -5a^2 \sin at \hat{i} + 2b \hat{j} \\
 \text{a) } \vec{L}|_{t=1} &= m \vec{v}|_{t=1} = 2 [5\pi \cos \pi + 2(0.25) \hat{j} + 2 \hat{k}] \\
 &= 2 [-15.708 \hat{i} + 0.5 \hat{j} + 2 \hat{k}] \\
 \vec{L}|_{t=1} &= -10\pi \hat{i} + \hat{j} + 4 \hat{k} \\
 \text{b) } \vec{F}|_{t=1} &= m \vec{a}|_{t=1} \\
 &= 2 [-5\pi^2 \sin \pi \hat{i} + 0.5 \hat{j}] \\
 \vec{F} &= 1 \hat{j} \text{ N}
 \end{aligned}$$

12.2.18 A particle A has mass m_A and velocity $\vec{v}_A$. A particle B at the same location has mass $m_B = 2m_A$ and velocity equal to the other $\vec{v}_B = \vec{v}_A$. Point C is a reference point. What is the relationship

between:

- a) $\vec{L}_A$ and $\vec{L}_B$,
- b) $\vec{H}_{A/C}$ and $\vec{H}_{B/C}$, and
- c) E_{KA} and E_{KB} ?

(37)

12.2.18

$$m_B = 2m_A$$

$$\vec{v}_B = \vec{v}_A$$

$$\vec{r}_B = \vec{r}_A$$

$$(i) \vec{L}_B = m_B \vec{v}_B = 2m_A \vec{v}_A = 2\vec{L}_A$$

$$\vec{L}_B = 2\vec{L}_A$$

$$(ii) \vec{H}_{B/C} = \vec{r}_{B/C} \times \vec{L}_B = \vec{r}_{A/C} \times 2\vec{L}_A$$

$$\vec{H}_{B/C} = 2\vec{H}_{A/C}$$

$$(iii) E_{KB} = \frac{1}{2} m_B \vec{v}_B \cdot \vec{v}_B = 2 \left(\frac{1}{2} m_A \vec{v}_A \cdot \vec{v}_A \right) = 2 E_{KA}$$

$$E_{KB} = 2 E_{KA}$$

12.2.19 A particle of mass $m = 3 \text{ kg}$ moves in space. Its position, velocity, and acceleration at a given time are $\vec{r} = 2 \text{ m}\hat{i} + 3 \text{ m}\hat{j} + 5 \text{ m}\hat{k}$, $\vec{v} = -3 \text{ m/s}\hat{i} + 8 \text{ m/s}\hat{j} + 10 \text{ m/s}\hat{k}$, and $\vec{a} = -5 \text{ m/s}^2\hat{i} + 12 \text{ m/s}^2\hat{j} + 16 \text{ m/s}^2\hat{k}$, respectively. For this particle at the instant of interest, find its:

- a) linear momentum $\vec{L}$,
b) rate of change of linear momen-

tum $\dot{\vec{L}}$,

- c) angular momentum about the origin $\vec{H}_{/O}$,
d) rate of change of angular momentum about the origin $\dot{\vec{H}}_{/O}$,
e) kinetic energy E_K , and
f) rate of change of kinetic energy $\dot{E}_K$.

38

11.2.19

$$m = 3 \text{ kg}$$

$$\begin{aligned} \text{a) } \vec{L} &= m\vec{v} \\ &= 3[-3\hat{i} + 8\hat{j} + 10\hat{k}] \text{ kg m/sec} \end{aligned}$$

$$\vec{L} = [-9\hat{i} + 24\hat{j} + 30\hat{k}] \text{ kg m/sec}$$

$$\begin{aligned} \text{b) } \dot{\vec{L}} &= m \frac{d\vec{v}}{dt} = m\vec{a} \\ &= 3[-5\hat{i} + 12\hat{j} + 16\hat{k}] \text{ kg m/sec}^2 \\ \dot{\vec{L}} &= [-15\hat{i} + 36\hat{j} + 48\hat{k}] \text{ kg m/sec}^2 \end{aligned}$$

$$\begin{aligned} \text{c) } \vec{H}_{/O} &= \vec{r}_{/O} \times m\vec{v} \\ &= [2\hat{i} + 3\hat{j} + 5\hat{k}] \times [-9\hat{i} + 24\hat{j} + 30\hat{k}] \text{ kg m}^2/\text{sec} \\ &= [(90 - 120)\hat{i} + (-60 - 45)\hat{j} + (48 + 27)\hat{k}] \text{ kg m}^2/\text{sec} \\ \vec{H}_{/O} &= [-30\hat{i} - 105\hat{j} + 75\hat{k}] \text{ kg m}^2/\text{sec} \end{aligned}$$

$$\begin{aligned} \text{d) } \dot{\vec{H}}_{/O} &= \vec{r}_{/O} \times m\vec{a} \\ &= [2\hat{i} + 3\hat{j} + 5\hat{k}] \times [-15\hat{i} + 36\hat{j} + 48\hat{k}] \text{ kg m}^2/\text{sec}^2 \\ \dot{\vec{H}}_{/O} &= [-36\hat{i} - 171\hat{j} + 117\hat{k}] \text{ kg m}^2/\text{sec}^2 \end{aligned}$$

$$\text{e) } E_K = \frac{1}{2} m \vec{v} \cdot \vec{v} = \frac{3}{2} [3^2 + 8^2 + 10^2] = 256.5 \text{ Joules}$$

$$\begin{aligned} \text{f) } \dot{E}_K &= m\vec{a} \cdot \vec{v} = [-15\hat{i} + 36\hat{j} + 48\hat{k}] \cdot [3\hat{i} + 8\hat{j} + 10\hat{k}] \\ &= 45 + 288 + 480 = 813 \text{ Joules/sec} \end{aligned}$$

#24 Solution Michelle deSouza

Monday, March 8, 2021 1:31 PM

12.2.19

Given:

- mass $m = 3 \text{ kg}$
- $\vec{r} = 2\hat{i} + 3\hat{j} + 5\hat{k} \text{ m}$
- $\vec{v} = -3\hat{i} + 8\hat{j} + 10\hat{k} \text{ m/s}$
- $\vec{a} = -5\hat{i} + 12\hat{j} + 16\hat{k} \text{ m/s}^2$

Solve at this instant in time for:

a) linear momentum $\vec{L}$

$$\begin{aligned}\vec{L} &= m\vec{v} \\ &= 3 \text{ kg} (-3\hat{i} + 8\hat{j} + 10\hat{k}) \text{ m/s} \\ \vec{L} &= -9\hat{i} + 24\hat{j} + 30\hat{k} \text{ kg}\cdot\text{m/s}\end{aligned}$$

b) rate of change of linear momentum $\dot{\vec{L}}$

$$\begin{aligned}\dot{\vec{L}} &= \frac{d}{dt}\vec{L} = m\vec{a} \\ &= 3 \text{ kg} (-5\hat{i} + 12\hat{j} + 16\hat{k}) \text{ m/s}^2 \\ \dot{\vec{L}} &= -15\hat{i} + 36\hat{j} + 48\hat{k} \text{ N} \quad 1 \frac{\text{kg}\cdot\text{m}}{\text{s}^2} = 1 \text{ N}\end{aligned}$$

c) angular momentum about origin $\vec{H}_O$

$$\begin{aligned}\vec{H}_O &= \vec{r} \times m\vec{v} \\ \vec{H}_O &= (2\hat{i} + 3\hat{j} + 5\hat{k}) \text{ m} \times 3 \text{ kg} (-3\hat{i} + 8\hat{j} + 10\hat{k}) \text{ m/s} \\ \vec{r} \times \vec{v} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 5 \\ -3 & 8 & 10 \end{vmatrix} \rightarrow \begin{vmatrix} 3 & 5 \\ 8 & 10 \end{vmatrix} \hat{i} - \begin{vmatrix} 2 & 5 \\ -3 & 10 \end{vmatrix} \hat{j} + \begin{vmatrix} 2 & 3 \\ -3 & 8 \end{vmatrix} \hat{k} \\ &= [3(10) - 8(5)]\hat{i} - [2(10) - (-3)(5)]\hat{j} + [2(8) - (-3)(3)]\hat{k} \\ \vec{r} \times \vec{v} &= -10\hat{i} - 35\hat{j} + 25\hat{k} \\ \vec{H}_O &= -30\hat{i} - 105\hat{j} + 75\hat{k} \text{ kg}\cdot\text{m}^2/\text{s}\end{aligned}$$

d) rate of change of angular momentum about origin $\dot{\vec{H}}_O$

$$\begin{aligned}\dot{\vec{H}}_O &= \vec{r} \times m\vec{a} \\ &= (2\hat{i} + 3\hat{j} + 5\hat{k}) \text{ m} \times 3 \text{ kg} (-5\hat{i} + 12\hat{j} + 16\hat{k}) \text{ m/s}^2 \\ \dot{\vec{H}}_O &= -36\hat{i} - 171\hat{j} + 117\hat{k} \text{ N}\cdot\text{m}\end{aligned}$$

e) kinetic energy E_k

$$\begin{aligned}E_k &= \frac{1}{2} m \vec{v} \cdot \vec{v} \\ &= \frac{1}{2} (3 \text{ kg}) (-3\hat{i} + 8\hat{j} + 10\hat{k}) \cdot (-3\hat{i} + 8\hat{j} + 10\hat{k}) \text{ m/s} \\ E_k &= 259.5 \text{ J} \quad 1 \text{ J} = 1 \frac{\text{kg}\cdot\text{m}^2}{\text{s}^2}\end{aligned}$$

f) rate of change of kinetic energy $\dot{E}_k$

$$\begin{aligned}\dot{E}_k &= \frac{d}{dt} \left(\frac{1}{2} m \vec{v} \cdot \vec{v} \right) \\ &= \frac{1}{2} (3 \text{ kg}) (-5\hat{i} + 12\hat{j} + 16\hat{k}) \cdot (-3\hat{i} + 8\hat{j} + 10\hat{k}) \text{ m/s} \\ &= 406.5 \text{ W} \quad 1 \text{ W} = 1 \text{ J/s}\end{aligned}$$

813

Derivations

$$\begin{aligned}\vec{F} &= m\vec{a} & \vec{a} &= \frac{d}{dt}(\vec{v}), m = \text{const.} \\ \vec{F} &= \frac{d}{dt}(m\vec{v}) & m\vec{v} &= \vec{L} \\ \vec{F} &= \dot{\vec{L}} & \vec{F} &= \text{force sum} \\ & & \vec{L} &= \text{linear momentum}\end{aligned}$$

$$\begin{aligned}\vec{F} &= m\vec{a} & \vec{r} \times \{ \} \\ \vec{r} \times \vec{F} &= \vec{r} \times m\vec{a} & \rightarrow \text{try } \frac{d}{dt}(\vec{r} \times m\vec{v}) \\ \vec{r} \times \vec{F} &= \frac{d}{dt}(\vec{r} \times m\vec{v}) & = \frac{d\vec{r}}{dt} \times m\vec{v} + \vec{r} \times m \frac{d\vec{v}}{dt} \\ \vec{M} &= \frac{d}{dt}(\vec{H}_O) & \vec{M} = \text{moment sum} \\ & & \vec{H}_O = \text{angular momentum} \\ & & = \vec{v} \times m\vec{v} + \vec{r} \times m\vec{a} \\ & & = \vec{r} \times m\vec{a} \checkmark\end{aligned}$$

only works if $\frac{d\vec{r}}{dt} = \vec{v}$ and $\frac{d\vec{v}}{dt} = \vec{a}$
aka $\vec{r}$, $\vec{v}$, and $\vec{a}$ are defined correctly
relative to each other — e.g. if $\vec{r}$ defined
relative to still point and $\vec{v}$ defined relative
to moving point, AMS does not work!

$$\begin{aligned}\vec{F} &= m\vec{a} & \{ \} \cdot \vec{v} \\ \vec{F} \cdot \vec{v} &= m\vec{a} \cdot \vec{v} & \rightarrow \text{try } \frac{d}{dt}(\frac{1}{2} m \vec{v} \cdot \vec{v}) \\ \vec{F} \cdot \vec{v} &= \frac{d}{dt}(\frac{1}{2} m \vec{v} \cdot \vec{v}) & \vec{v} \cdot \vec{v} = v^2 \\ P &= \frac{d}{dt}(\frac{1}{2} m v^2) & P = \text{power} \\ P &= \dot{E}_k & E_k = \text{kinetic energy} \\ & & = \frac{1}{2} m \vec{a} \cdot \vec{v} + \frac{1}{2} m \vec{v} \cdot \vec{a} \\ & & = m \vec{a} \cdot \vec{v} \checkmark\end{aligned}$$

Similar warning applies! E.g. if $\vec{a}$ is defined
relative to an accelerating point, this derivation
does not work

12.2.20 A particle has position $\vec{r} = 3\text{ m}\hat{i} - 2\text{ m}\hat{j} + 4\text{ m}\hat{k}$, velocity $\vec{v} = 2\text{ m/s}\hat{i} - 3\text{ m/s}\hat{j} + 7\text{ m/s}\hat{k}$, and acceleration $\vec{a} = 1\text{ m/s}^2\hat{i} - 8\text{ m/s}^2\hat{j} + 3\text{ m/s}^2\hat{k}$. For each position of a point P defined below, find the rate of change of angular momentum, $\dot{\vec{H}}_P$, of the particle with respect to

the point P .

- $\vec{r}_P = 3\text{ m}\hat{i} - 2\text{ m}\hat{j} + 4\text{ m}\hat{k}$,
- $\vec{r}_P = 6\text{ m}\hat{i} - 4\text{ m}\hat{j} + 8\text{ m}\hat{k}$,
- $\vec{r}_P = -9\text{ m}\hat{i} + 6\text{ m}\hat{j} - 12\text{ m}\hat{k}$, and
- $\vec{r}_P = \vec{0}$

(39)

11.2.20

$$\begin{aligned}\vec{H}_P &= \vec{r}_P \times m\vec{a} \\ &= (\vec{r} - \vec{r}_P) \times m\vec{a}\end{aligned}$$

$$\begin{aligned}\text{a) } \vec{H}_P &= m \left\{ (3\hat{i} - 2\hat{j} + 4\hat{k}) - (3\hat{i} - 2\hat{j} + 4\hat{k}) \right\} \times [\hat{i} - 8\hat{j} + 3\hat{k}] \\ &= m [0 \times (\hat{i} - 8\hat{j} + 3\hat{k})] \\ \dot{\vec{H}}_P &= 0\end{aligned}$$

$$\begin{aligned}\text{b) } \vec{H}_P &= m \left[(3\hat{i} - 2\hat{j} + 4\hat{k}) - (6\hat{i} - 4\hat{j} + 8\hat{k}) \right] \times [\hat{i} - 8\hat{j} + 3\hat{k}] \\ &= -m [(3\hat{i} - 2\hat{j} + 4\hat{k}) \times (\hat{i} - 8\hat{j} + 3\hat{k})] \\ \dot{\vec{H}}_P &= -m [26\hat{i} - 5\hat{j} - 22\hat{k}]\end{aligned}$$

$$\begin{aligned}\text{c) } \vec{H}_P &= m \left[(3\hat{i} - 2\hat{j} + 4\hat{k}) - (-9\hat{i} + 6\hat{j} - 12\hat{k}) \right] \times [\hat{i} - 8\hat{j} + 3\hat{k}] \\ &= m [12\hat{i} - 8\hat{j} + 16\hat{k}] \times [\hat{i} - 8\hat{j} + 3\hat{k}] \\ \dot{\vec{H}}_P &= m [104\hat{i} - 20\hat{j} - 92\hat{k}]\end{aligned}$$


12.2.21 A particle of mass $m = 6 \text{ kg}$ is moving in space. Its position, velocity, and acceleration at some instant are $\vec{r} = 1 \text{ m}\hat{i} - 2 \text{ m}\hat{j} + 4 \text{ m}\hat{k}$, $\vec{v} = 3 \text{ m/s}\hat{i} + 4 \text{ m/s}\hat{j} - 7 \text{ m/s}\hat{k}$, and $\vec{a} = 5 \text{ m/s}^2\hat{i} + 11 \text{ m/s}^2\hat{j} - 9 \text{ m/s}^2\hat{k}$, respectively. At this instant, find:

- the net force $\sum \vec{F}$ on the particle,
- the net moment on the particle about the origin $\sum \vec{M}_O$ due to the

applied forces, and

- the power P of the applied forces.

Particle FBD



Problem 12.21: FBD of the particle

11.2.22

$$\begin{aligned} \text{a)} \quad \sum \vec{F} &= \vec{L} = m\vec{a} \\ &= 6 [5\hat{i} + 11\hat{j} - 9\hat{k}] \\ \vec{F} &= 30\hat{i} + 66\hat{j} - 54\hat{k} \text{ N} \end{aligned}$$



$$\begin{aligned} \text{b)} \quad \sum \vec{M}_O &= \vec{H}_O \\ &= m \vec{r}_O \times \vec{a}_O \\ &= m [\hat{i} - 2\hat{j} + 4\hat{k}] \times [5\hat{i} + 11\hat{j} - 9\hat{k}] \\ &= 6 [-26\hat{i} + 29\hat{j} + 21\hat{k}] \\ \sum \vec{M}_O &= -156\hat{i} + 174\hat{j} + 126\hat{k} \text{ N}\cdot\text{m} \end{aligned}$$

$$\begin{aligned} \text{c) Power,} \quad P &= \sum m\vec{a} \cdot \vec{v} \\ &= 6 [5\hat{i} + 11\hat{j} - 9\hat{k}] \cdot [3\hat{i} + 4\hat{j} - 7\hat{k}] \\ &= 6 [15 + 44 - 63] \\ P &= -24 \text{ Joules/sec} \end{aligned}$$

12.2.22 At a time of interest, a particle with mass $m_1 = 5 \text{ kg}$ has position, velocity, and acceleration $\vec{r}_1 = 3 \text{ m}\hat{i}$, $\vec{v}_1 = -4 \text{ m/s}\hat{j}$, and $\vec{a}_1 = 6 \text{ m/s}^2\hat{j}$, respectively. Another particle with mass $m_2 = 5 \text{ kg}$ has position, velocity, and acceleration $\vec{r}_2 = -6 \text{ m}\hat{i}$, $\vec{v}_2 = 5 \text{ m/s}\hat{j}$, and $\vec{a}_2 = -4 \text{ m/s}^2\hat{j}$, respectively. For this system of two particles, and at this time, find its

a) linear momentum $\vec{L}$,

- b) rate of change of linear momentum $\dot{\vec{L}}$
 c) angular momentum about the origin $\vec{H}_O$,
 d) rate of change of angular momentum about the origin $\dot{\vec{H}}_O$,
 e) kinetic energy E_K , and
 f) rate of change of kinetic energy $\dot{E}_K$.

40

11-2-21

$$\begin{array}{l|l} m_1 = 5 \text{ kg} & m_2 = 5 \text{ kg} \\ \vec{r}_1 = 3\hat{i} \text{ m} & \vec{r}_2 = -6\hat{i} \\ \vec{v}_1 = -4\hat{j} \text{ m/s} & \vec{v}_2 = 5\hat{j} \\ \vec{a}_1 = 6\hat{j} \text{ m/s}^2 & \vec{a}_2 = -4\hat{j} \end{array}$$

$$\begin{aligned} \text{a) } \vec{L} &= \sum m_i \vec{v}_i \\ &= 5(-4\hat{j}) + 5(5\hat{j}) \\ \vec{L} &= 5\hat{j} \text{ kg m/sec} \end{aligned}$$

$$\begin{aligned} \text{b) } \dot{\vec{L}} &= \sum m_i \vec{a}_i \\ &= 5(6\hat{j}) + 5(-4\hat{j}) \\ \dot{\vec{L}} &= 10\hat{j} \text{ kg m/sec}^2 \end{aligned}$$

$$\begin{aligned} \text{c) } \vec{H}_O &= \sum \vec{r}_{i/O} \times m\vec{v}_i \\ &= 5[3\hat{i} \times (-4\hat{j}) + (-6\hat{i}) \times (5\hat{j})] \\ \vec{H}_O &= -210\hat{k} \text{ kg m}^2/\text{sec}^2 \end{aligned}$$

$$\begin{aligned} \text{d) } \dot{\vec{H}}_O &= \sum \vec{r}_{i/O} \times m\vec{a}_i \\ &= 5[3\hat{i} \times 6\hat{j} + (-6\hat{i}) \times (-4\hat{j})] \\ &= 5[18\hat{k} + 24\hat{k}] \\ \dot{\vec{H}}_O &= 210\hat{k} \text{ kg m}^2/\text{sec}^2 \end{aligned}$$

$$\text{e) } E_K = \sum \frac{1}{2} m_i \vec{v}_i^2 = \frac{5}{2} [(4)^2 + (5)^2] = 102.5 \text{ Joules}$$

$$E_K = 102.5 \text{ Joules}$$

$$\begin{aligned} f) \dot{E}_K &= \sum m \vec{a} \cdot \vec{v} \\ &= 5 [(6\hat{j}) \cdot (-4\hat{j}) + (-4\hat{j}) \cdot (5\hat{j})] \\ &= 5 [-24 - 20] \\ \dot{E}_K &= -220 \text{ Joules/sec} \end{aligned}$$

12.2.23 A particle of mass $m = 250$ gm is shot straight up (parallel to the y -axis) from the x -axis at a distance $d = 2$ m from the origin. The velocity of the particle is given by $\vec{v} = v\hat{j}$ where $v^2 = v_0^2 - 2ah$, $v_0 = 100$ m/s, $a = 10$ m/s² and h is the height of the particle from the x -axis.

- a) Find the linear momentum of the particle at the outset of motion ($h = 0$).

- b) Find the angular momentum of the particle about the origin at the outset of motion ($h = 0$).
- c) Find the linear momentum of the particle when the particle is 20 m above the x -axis.
- d) Find the angular momentum of the particle about the origin when the particle is 20 m above the x -axis.

(42)

$$m = 250 \text{ grams} = \frac{1}{4} \text{ kg}$$

$$\vec{r}_0 = 2\hat{i}$$

$$\vec{v} = \sqrt{v_0^2 - 2ah}\hat{j}$$

$$v_0 = 100 \text{ m/sec}$$

$$a = 10 \text{ m/sec}^2$$

$$h = y \quad \text{if } \vec{r}_{p/o} = x\hat{i} + y\hat{j}$$

$$a) \quad \vec{L}|_{t=0} = m\vec{v}|_{t=0}$$

$$= \frac{1}{4} \left[\sqrt{v_0^2 - 2a(0)} \hat{j} \right] \text{ kg m/sec}$$

$$= \frac{v_0}{4} \hat{j} \text{ kg m/sec}$$

$$\vec{L}|_{t=0} = 25 \hat{j} \text{ kg m/sec}$$

$$b) \quad \vec{H}|_{t=0} = \vec{r}_0 \times m\vec{v}|_{t=0}$$

$$= 2\hat{i} \times 25\hat{j}$$

$$\vec{H}|_{t=0} = 50 \hat{k} \text{ kg m}^2/\text{sec}$$

$$c) \quad \text{for } h = 20 \text{ m}$$

$$\vec{L} = m\vec{v}$$

$$= \frac{1}{4} \left[\sqrt{v_0^2 - 2ah} \right] \hat{j}$$

$$\vec{L} = 24.5 \hat{j} \text{ kg m/sec}$$

$$d) \quad \text{for } h = 20 \text{ m}$$

$$\vec{r}_{p/o} = \hat{i} + 20\hat{j}$$

$$\vec{H}_{p/o} = \vec{r}_{p/o} \times m\vec{v}$$

$$= (2\hat{i} + 20\hat{j}) \times (24.5\hat{j})$$

$$\vec{H}_{p/o} = 99 \hat{k} \text{ kg m}^2/\text{sec}$$

Disclaimer: We have lost track of the many people who wrote these solutions. They are not official and have not been reviewed by the authors for correctness.